NAME	 	 	
ID#			

ORGANIC CHEMISTRY I (2301)

9:30 – 10:20 am, June 30, 2015

Exam 1

If you want to pick this exam up on Thursday in class (in public), please check the box on the right:	
If you do not check the box, I will not bring your exam to class on Thursday, and you will need to pick up your exam in private from Chemistry department staff in 115 Smith beginning Thursday afternoon. Exa not picked up within two weeks will be disposed of.	ms that are

A periodic table is attached to the back of this exam as an aid. Otherwise, you are not permitted to use any other materials (including notes, books, or electronic devices of any kind).

Right now, write your name and student ID number at the top of this page. When the exam begins, please write your name at the top of the next page.

You may use pen or pencil. However, re-grades will be considered only for exams completed in pen.

Please write your answers in the boxes/spaces provided. If your answer is not in the appropriate space (say, for example, it's on the back of the page), draw us an arrow and/or note telling us where to look.

NAME				

1. /6 5. /18 Scoring:

2. / 20

6. / 20


3. / 12

7. / 6

4. _____/ 18


Total Score: _____/ 100

1. (6 pts) Draw Lewis dash-bond structures for two constitutional isomers that have molecular formula CH₄N₂, and that have no formal charges on any atom. Draw all atoms and lone pairs of electrons.





2. (20 pts) For each of the molecules on the left, draw as many of the best Lewis dash-bond resonance structures as there are boxes to put them in. (Feel free to omit lone pairs and C-H's, or draw them—your choice.) Then, below each resonance structure, describe whether each would be a major or minor contributor. Finally, draw a resonance hybrid that illustrates partial charges and multiple bonds.

resonance structure resonance hybrid $H_2^{\bigcirc}N_{\bigcirc}O$ contribution: contribution: (major/minor?)



and/or torsional strain? Circle your answers in the boxes below. Then, if you circled "YES" in either box, illustrate the electron-pair repulsion (the strain) on my skeleton structure with a double-headed arrow. You may have to add to my drawing in order to show the repelling electrons. (If you circled "NO", leave the rest of that box blank.)








b. 1,4-Dioxane puckers from planarity in order to avoid angle and torsional strain. In the box on the right, draw the most stable conformation of 1,4-dioxane. In your drawing, **include all H atoms**, but feel free to omit lone pairs.




- 4. (18 pts) For each of the Lewis structures drawn below, in the boxes provided:
 - Draw Lewis wedge/dashed-bond structures that illustrate the most stable three-dimensional structure of the molecule. Draw all atoms, but feel free to omit lone pairs.
 - In the boxes provided, write the hybridization state for any atom heavier than hydrogen.
 - In the boxes provided, give any bond angle indicated by curved arrows in the original Lewis structure.











- 5. (18 pts) For each of the sets of molecules below:
 - Using "electron pushing" (with double-barbed arrows), show how the molecules on the left would react in an acid-base reaction to transfer a proton from one to the other.
 - In the box on the right, draw the conjugate acid and base products of each reaction.
 - In the middle, draw an equilibrium arrow that shows whether you feel the acid-base equilibrium would lie on the left or the right.



6. (20 pts)

a. For the difluorobutane shown at right, in the boxes below, draw Newman projections that show the most stable, second-most stable, least stable, and second-least stable conformations of the molecule. Draw your projections looking down the central C-C bond, using the perspective I've shown in the drawing.



b. Different conformations can contribute different polarities to the overall, average polarity of a molecule. For each of the two most stable conformations of the molecule above, circle whether the molecule is polar or non-polar when it is in that conformation. If you circle "POLAR", also draw one dipole arrow (+>>) that shows the total dipole moment for that conformer.

Newman projection for most stable conformation	Newman projection for second-most stable conformation
POLAR or NON-POLAR ?	POLAR or NON-POLAR ?
Newman projection for least stable conformation	Newman projection for second-least stable conformation

c. Would you call the 2,3-difluorobutene shown at right a


CONFIGURATIONAL ISOMER,

STEREOISOMER, o


r NEITHER

compared to the 2,3-difluorobutane on the previous page? (Circle one answer.)

d. Which is more polar:



7. (6 pts) Each of the basic molecules below has multiple potential protonation sites. Given the pKa values in the chart on the right, draw the organic product you would expect if each molecule was combined with just one molecule of H₃O⁺.



18 8A 2 He lium 4.00	10 Neon 20.18	18 Ar	39.95	98 2	Krypton 83.80	24 X	Xenon 131.29	% Z	Radon (222)			71
17 7A	9 Fluorine 19.00	17 C Ploying	35.45	35 પૂ	Bromine 79.90	- 23	lodine 126.90	85 At	Astatine (210)			70
16 6A	0,											69
15 5A	7 N Nitrogen 14.01	15 P	30.97	33 Ac	Arsenic 74.92	51 Sb	Antimony 121.76	88 	Bismuth 208.98			89
4 4 4 A	6 Carbon 12.01	44 S	28.09	32 G	Germanium 72.61	20 S	Tin 118.71	85 Pb	Lead 207.2			67
13 3A						I						99
		- 6	2B	% K	Zinc 65.39	84 C	Cadmium 112.41	8 5	Mercury 200.59			65
		Ξ	1 1B	5 ₂	Copper 63.55	47 A q	Silver 107.87	79 Au	Gold 196.97			64
		Ç	2	58 Z	Nickel 58.69	46 Pd	Palladium 106.42	8 ⊈	Platinum 195.08			63
	oer Ibol	nic mass*	- 1 1								(268)	69
Key	Atomic number Element symbol Element name	Average atomic mass	, [۵	Iron 55.85	44 B C	Ruthenium 101.07	92 Os	Osmium 190.23	108 HS	(269)	61
*	+++	_	, 7B	25 Mn	ž	43 C	ĕ	75 Be	Rhenium 186.21	107 Bh	(264)	09
	11 — Sodium -	5. J	éB	5 5	⊑ _	42 Mo	Molybdenum 95.94	74 X	Tungsten 183.84	106 Sg	(266)	29
		ער	5B	23	Vanadium 50.94	4 Z	Niobium 92.91	73 a	Fantalum 180.95	105 Db	(262)	28
		4	4B	Z F		40 Z	Zirconium 91.22	₹ 5	Hafnium 178.49	104 Pf	(261)	
		ď	3B	7 م	Scandium 44.96	% ➤	Yttrium 88.91	57 Ea	Lanthanum 138.91	89 Ac	(227)	
2A	Be Beryllium 9.01	Mg Mg	24.31	02 C	Calcium 40.08	% %	Strontium 87.62	56 Ba	Barium 137.33	88 Ba		
1 1A 1 Hydrogen 1.01	3 Lithium 6.94		22.99	€ 🛪	Potassium 39.10	37 B	Rubidium 85.47	55 Cs	Cesium 132.91	87 Fr	(223)	
-	α	က		•	4	L	n	U	0	7		

* If this number is in parentheses, then it refers to the atomic mass of the most stable isotope.

28	29	09	61	62	63	64	65	99	29	89	69	20	71
ပီ	P	PZ	Pm	Sm	Eu	<u> </u>	Д	٥	운	щ	٦	Υb	ב
Cerium	Praseodymium	Neodymium	Promethium	Samarium	Europium	Gadolinium	Terbium	Dysprosium	Holminm	Erbium	Thulium	Ytterbium	Lutetium
140.12	140.91	144.24	(145)	150.36	151.96	157.25	158.93	162.50	164.93	167.26	168.93	173.04	174.97
06	91	92	63	94	92	96	26	86	66	100	101	102	103
드	Ра	>	dN	Pu	Am	CH	BK	℧	Es	FB	Md	٥ N	۲
Thorium	Protactinium		Neptunium	Plutonium	Americium	Curium	Berkelium	Californium	Einsteinium	Fermium	Mendelevium	Nobelium	Lawrencium
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(262)

Copyright © 2008 California Department of Education