NAME	 	 	
ID#			

ORGANIC CHEMISTRY I (2301)

9:30 - 10:20 am, July 14, 2015

Exam 2

If you want to pick this exam up on Thursday in class (in public), please check the box on the right:	
If you do not check the box, I will not bring your exam to class on Tuesday, and you will need to pick up your exam in private from Chemistry department staff in 115 Smith beginning Friday, July 17 th . Exams picked up within two weeks will be disposed of.	that are not

A periodic table is attached to the back of this exam as an aid. Otherwise, you are not permitted to use any other materials (including notes, books, or electronic devices of any kind).

Right now, write your name and student ID number at the top of this page. When the exam begins, please write your name at the top of the next page.

You may use pen or pencil. However, re-grades will be considered only for exams completed in pen.

Please write your answers in the boxes/spaces provided. If your answer is not in the appropriate space (say, for example, it's on the back of the page), draw us an arrow and/or note telling us where to look.

NAME						

Scoring:

Total Score: / 100

1. (12 pts) How would you describe the relationship between each of the pairs of structures below? Are they enantiomers or diastereomers, or are they just two ways of illustrating the same molecule? Circle one answer for each pair.

$$CI \xrightarrow{CH_3} CH_3$$

$$\begin{array}{c} H \\ \downarrow \\ H \\ CH_3 \end{array} CI$$

ENANTIOMERS

or

DIASTEREOMERS

or

SAME MOLECULE

ENANTIOMERS

or

DIASTEREOMERS

or

SAME MOLECULE

ENANTIOMERS


or

DIASTEREOMERS


or

SAME MOLECULE

2. (16 pts) On the structures below, **label each chiral center** with its appropriate Cahn-Ingold-Prelog designation [(R) or (S)]. Make it clear which atom in the drawing you are labeling. Then, for each structure, **circle** whether you think the molecule is chiral or achiral.



3. (20 pts) Draw the missing reactant or product in the empty boxes. For products, give the predominant, most favored product. Illustrate stereochemistry in your answer where appropriate. For reactions that yield multiple enantiomers, draw only one enantiomer in the box, and include the note "+ enantiomer".



$$CH_3O^{\bigodot}Na^{\bigodot}$$

$$an \ alkyl \ halide$$

$$H_3C \xrightarrow{CH_3} CH_3$$

$$(as \ the \ major \ product)$$

$$CH_3 \cap CH_3$$

$$(as \ the \ major \ product)$$

4. (11 pts) For each reaction shown below, **circle all possible products** (including minor ones). Keep in mind that, for each reaction, you might circle one, multiple, or no products.

$$\begin{array}{c|c}
& & & \\
& & \\
\hline
\hline
CH_3
\end{array}$$


$$\begin{array}{c}
& & \\
& \\
\hline
CH_3
\end{array}$$

$$\begin{array}{c}
& \\
& \\
& \\
\hline
CH_3
\end{array}$$

$$\begin{array}{c}
& \\
& \\
& \\
\hline
CH_3
\end{array}$$


5. (26 pts) Ammonia reacts with *tert*-butylchloride to yield two sets of products:

a. The two products are made from the same rate-determining transition state, and through the same reactive intermediate. What are the structures of those two states?



- b. On the diagram on the next page, draw potential energy curves that describe the formation of each of your two products. For each curve:
 - Draw energy levels for all transition states and intermediates, and connect them with curves. You do not need to draw any chemical structures, just energies.
 - Energy levels for the starting material and the two products are given.
 - Wherever your two paths share common intermediates or transition states, you only need to draw one curve; draw two separate curves only where the paths diverge.

starting _____



How would the reaction change if ammonia was replaced with diisopropylethylamine (DIPEA)?

Using DIPEA would make the overall reaction

FASTER SLOWER SAME RATE

Using DIPEA would make the product ratio A/B

INCREASE DECREASE STAY THE SAME (favor A over B) (favor B over A)

- 6. (15 pts) For the reaction shown below, draw a mechanism that explains how the product is generated from the starting material. In your answer, make sure that you:
 - Draw each step of the mechanism separately;
 - Use "electron pushing" to show where the electrons in each step go;
 - Use only the molecules that you are given; do not invoke reactants or solvents that aren't in the problem.

I have drawn the first set of electron-pushing arrows for you.

HO OH
$$H_2SO_4$$
 O

$$H - OSO_3H$$
 $HO \longrightarrow OH$

18 8A 2 Te Helium 4.00	10 Neon 20.18	18 Argon	39.95	 % ∑	Krypton 83.80	24	Xenon 131.29	% 2	Radon (222)			71
17 A 7	9 Fluorine 19.00	17 C	35.45	સ્ જે	Bromine 79.90	- 23	lodine 126.90	85 At	Astatine (210)			70
16 6A	0,											69
15 5A	7 N itrogen 14.01	15 P Phosphorus	30.97	33 As	Arsenic 74.92	51 5	Antimony 121.76	83 	Bismuth 208.98			89
4 4 4 A	6 Carbon 12.01	14 S	28.09	Ge 35	Germanium 72.61	20	118.71	85 Pb	Lead 207.2			67
3A 3A	5 B Boron 10.81	13 Al Aluminum	26.98	31 Ga	Gallium 69.72	49 2	Indium 114.82	= 84	Thallium 204.38			99
		12	2B	® Z	Zinc 65.39	8 ⁴ C	Cadmium 112.41	8 2	Mercury 200.59			65
		Ξ	1 18	ور 20	Copper 63.55	74 <	Silver 107.87	79 Au	Gold 196.97			64
		10		8 Z	Nickel 58.69	46 D	Palladium 106.42	82 T	Platinum 195.08			63
	oer Ibol ne	nic mass*	I L								Meitnerium (268)	69
Key	Atomic number Element symbol	Average atomic mass 8		56 Te	Iron 55.85	44	Ruthenium 101.07	92 Os	Osmium 190.23	108 Hs	Hassium (269)	61
<u> </u>	+++	 	7B	25 Mn	Š	£ 4 ⊢	<u>ĕ</u>		Rhenium 186.21	107 Bh	Bohrium (264)	9
	11— Sodium -	9	6B	გ ე	Chromium 52.00	45	Molybdenum 95.94	⁷ 8	Tungsten 183.84	106 Sg	Seaborgium (266)	29
		rO	5B	~ 53	Vanadium 50.94	4 Z	Niobium 92.91	2 23	Tantalum 180.95	105 Db	Dubnium (262)	228
		4		Z ;		4 ,	Zirconium 91.22	2 2	Hafnium 178.49	² ₹	Rutherfordium (261)	
			38	Sc	Scandium 44.96	e >	Yttrium 88.91	57 La	Lanthanum 138.91	88 Ac	Actinium (227)	
2 S	Be Beryllium	12 Mg	24.31	Ca 50	Calcium 40.08	ڻ %	SO .		Barium 137.33	88 Ba		
14 14 Hydrogen 1.01	3 Lithium 6.94	Nodiun Sodiun	22.99	<u>ნ</u> 🕶	Potassium 39.10	•	ш_				Francium (223)	
-	N	က		4			വ	Q	0			

* If this number is in parentheses, then it refers to the atomic mass of the most stable isotope.

28	29	09	61	62	63	64	92	99	29	89	69	20	71
ပီ	P	βN	Pm	Sm	Eu	<u> </u>	P	٥	운	щ	٦	Υb	ב
Cerium	Praseodymium	Neodymium	Promethium	Samarium	Europium	Gadolinium	Terbium	Dysprosium	Holminm	Erbium	Thulium	Ytterbium	Lutetium
140.12	140.91	144.24	(145)	150.36	151.96	157.25	158.93	162.50	164.93	167.26	168.93	173.04	174.97
06	91	62	63	94	92	96	26	86	66	100	101	102	103
드	Ра	-	dN	Pu	Am	CH	路	℧	Es	FB	Md	٥ N	۲
Thorium	Protactinium		Neptunium	Plutonium	Americium	Curium	Berkelium	Californium	Einsteinium	Fermium	Mendelevium	Nobelium	Lawrencium
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(262)

Copyright © 2008 California Department of Education