



Workshop 23 Solutions Complex Splitting Patterns in NMR

When we covered complex splitting patterns in lecture, I showed the simplest complex splitting pattern (if you will), the doublet of doublets; I also showed that the two coupling constants J_1 and J_2 that give rise to this pattern can be calculated by measuring the distance between peaks 1 and 2 (for J_1) and between peaks 1 and 3 (for J_2). There are three such patterns in this spectrum, corresponding to the three protons on the left-hand side of the molecule.




a. So, for each of those multiplets,




$$J_1: \Delta\delta = (7.2715 \text{ ppm}) - (7.2249 \text{ ppm}) \\ = 0.0466 \text{ ppm} \\ J_1 (\text{Hz}) = (0.0466 \text{ ppm}) \times (300 \text{ MHz}) \\ = 14.0 \text{ Hz}$$

$$J_2: \Delta\delta = (7.2715 \text{ ppm}) - (7.2507 \text{ ppm}) \\ = 0.0208 \text{ ppm} \\ J_2 (\text{Hz}) = (0.0208 \text{ ppm}) \times (300 \text{ MHz}) \\ = 6.2 \text{ Hz}$$



$$J_1: \Delta\delta = (4.8851 \text{ ppm}) - (4.8385 \text{ ppm}) \\ = 0.0466 \text{ ppm} \\ J_1 (\text{Hz}) = (0.0466 \text{ ppm}) \times (300 \text{ MHz}) \\ = 14.0 \text{ Hz}$$

$$J_2: \Delta\delta = (4.8851 \text{ ppm}) - (4.8802 \text{ ppm}) \\ = 0.0049 \text{ ppm} \\ J_2 (\text{Hz}) = (0.0049 \text{ ppm}) \times (300 \text{ MHz}) \\ = 1.5 \text{ Hz}$$



$$J_1: \Delta\delta = (4.5615 \text{ ppm}) - (4.5407 \text{ ppm})$$

$$= 0.0208 \text{ ppm}$$

$$J_1 (\text{Hz}) = (0.0208 \text{ ppm}) \times (300 \text{ MHz})$$

$$= 6.2 \text{ Hz}$$


$$J_2: \Delta\delta = (4.5615 \text{ ppm}) - (4.5566 \text{ ppm})$$

$$= 0.0049 \text{ ppm}$$


$$J_1 (\text{Hz}) = (0.0208 \text{ ppm}) \times (300 \text{ MHz})$$

$$= 1.5 \text{ Hz}$$


Importantly, each of the coupling constants in each multiplet matches one in another multiplet. That's because coupling constants between two nuclei are the same in both directions. So that means the three peaks represent three protons coupled to each other like



- b. So then which proton in vinyl acetate is which? Smith shows that J values of double-bonded protons look like this:



These values are pretty close to the ones measured in the multiplets. (Yes, 6.2 Hz is quite a bit less than 10 Hz, but the trend is there.) So, the full NMR assignment would be:



$$J_{ab} = 14.0 \text{ Hz}$$

$$J_{bc} = 1.5 \text{ Hz}$$

$$J_{ac} = 6.2 \text{ Hz}$$