

# Naming Alkanes and Alkyl Groups

**Table 2.1 Nomenclature and Physical Properties of Straight-Chain Alkanes**

Number of carbons	Molecular formula	Name	Condensed structure
1	CH <sub>4</sub>	methane	CH <sub>4</sub>
2	C <sub>2</sub> H <sub>6</sub>	ethane	CH <sub>3</sub> CH <sub>3</sub>
3	C <sub>3</sub> H <sub>8</sub>	propane	CH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub>
4	C <sub>4</sub> H <sub>10</sub>	butane	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>
5	C <sub>5</sub> H <sub>12</sub>	pentane	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>
6	C <sub>6</sub> H <sub>14</sub>	hexane	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub> CH <sub>3</sub>
7	C <sub>7</sub> H <sub>16</sub>	heptane	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CH <sub>3</sub>
8	C <sub>8</sub> H <sub>18</sub>	octane	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>6</sub> CH <sub>3</sub>
⋮	⋮	⋮	⋮

(linear alkanes are also called *n*-alkanes, where *n* is "normal")

Names of alkyl fragments derive from alkanes:

CH<sub>3</sub>—  
a methyl group

CH<sub>3</sub>CH<sub>2</sub>—  
an ethyl group

CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>—  
a propyl group

CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>—  
a butyl group


CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>—  
a pentyl group

R—  
any alkyl group


## Naming Functionalized Alkanes

CH <sub>4</sub> <i>methane</i>	CH <sub>3</sub> — (or H <sub>3</sub> C—) <i>methyl</i>	—OH <i>alcohol</i>	CH <sub>3</sub> —OH <i>methyl alcohol</i> ( <i>methanol</i> )
CH <sub>3</sub> CH <sub>3</sub> <i>ethane</i>	CH <sub>3</sub> CH <sub>2</sub> — <i>ethyl</i>	—O—C(=O)—R <i>ester</i>	CH <sub>3</sub> CH <sub>2</sub> —O—C(=O)—CH <sub>3</sub> <i>acetic acid ethyl ester</i> ( <i>ethyl acetate</i> )
CH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub> <i>propane</i>	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> — <i>n-propyl</i> <div style="display: inline-block; vertical-align: middle;"> <math>\begin{array}{c} \text{CH}_3 \\   \\ \text{CH} - \\   \\ \text{CH}_3 \end{array}</math> </div> <i>iso-propyl</i>	—NH <sub>2</sub> <i>amine</i>	<div style="display: inline-block; vertical-align: middle;"> <math>\begin{array}{c} \text{CH}_3 \\   \\ \text{CH} - \text{NH}_2 \\   \\ \text{CH}_3 \end{array}</math> </div> <i>isopropyl amine</i>
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub> <i>butane</i>	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> — <i>n-butyl</i> <div style="display: inline-block; vertical-align: middle;"> <math>\begin{array}{c} \text{CH}_3 \\   \\ \text{CH} - \\   \\ \text{CH}_2 \\   \\ \text{CH}_3 \end{array}</math> </div> <i>sec-butyl</i> <div style="display: inline-block; vertical-align: middle;"> <math>\begin{array}{c} \text{CH}_3 \\   \\ \text{C} - \\ / \quad \backslash \\ \text{CH}_3 \quad \text{CH}_3 \end{array}</math> </div> <i>tert-butyl</i>	—O— <i>ether</i>	<div style="display: inline-block; vertical-align: middle;"> <math>\begin{array}{c} \text{CH}_3 \\   \\ \text{H}_3\text{C} - \text{O} - \text{C} - \text{CH}_3 \\   \\ \text{CH}_3 \end{array}</math> </div> <i>methyl tert-butyl ether (MTBE)</i>


# Naming Functionalized Alkanes




*has format:*




**Primary** carbon, forms primary ester.




*has format:*



**Secondary** carbon, forms secondary amine.






*has format:*



**Tertiary** carbon, forms tertiary ether.


Just **methyl**, not primary.

# Bond Rotation and Newman Projections




# Rotation About Single Bond in Ethane


Newman projections:




“eclipsed”



“staggered”




“skew”



In “eclipsed” conformation, electron-pair repulsion between  $\sigma$  bonds is maximized.



# Rotation About Single Bond in Ethane



*Torsional* energy of 2.9 kcal/mol must be overcome to rotate 120°.

Timescale of that happening: once every 100 femtoseconds (at room temp).



## Rotation About Single Bond in Ethane



Video: Ethane, 273 K, 500 fs

(Simulation performed by molecular dynamics, using standard rotational and vibrational potentials.)


## Rotation About Single Bond in Ethane




Video: Ethane, 273 K, 500 fs

(Simulation performed by molecular dynamics, using standard rotational and vibrational potentials.)





# Rotation About Single Bond in Propane



# Rotation About Single Bond in Propane




## Rotation About Single Bond in Butane




*H-H steric interaction is unavoidable.*

*So, "totally eclipsed" is higher in energy than "eclipsed".*

## Rotation About Single Bond in Butane




## Rotation About Single Bond in Butane



### Points to note:




Gauche is not that much less stable than anti; so, butane spends plenty of time in gauche conformation.

Barrier to rotation of end methyl groups (3.3 kcal/mol) is less than that of internal C-C bond (5 kcal/mol).



## Conformational Preferences Extend to Higher Alkanes


*For linear alkanes, the all-anti conformation is preferred...*



*What Newman projections would you draw for these two perspectives?*

## Conformational Preferences Extend to Higher Alkanes

*For linear alkanes, the all-anti conformation is preferred...*



*...but not by much. Gauche conformers frequently observed.*